

# Leek High School



## Infection Control Policy

### Important Coronavirus (COVID-19) Update

This policy has been updated in line with the current guidance from the UK government. We have included an amendment to the policy in [Appendix A](#) which provides detailed infection control considerations for the current circumstances. Please have due regard for the stipulations within this amendment for the duration of schools being partially closed. Once students return to school full time, schools can refer to the provisions within the main body of this policy.

The DfE coronavirus helpline is: 0800 046 8687 Monday to Friday from 8:00am to 6:00pm and weekends 10:00am to 4:00pm. You can also email the helpline on:

[DfE.coronavirushelpline@education.gov.uk](mailto:DfE.coronavirushelpline@education.gov.uk)

This policy is available on our school website and is available on request from the school office.

This policy will be reviewed in full by the Local Governing Body on an annual basis. This policy is due for review in October 2021.

Headteacher

Date:

21-10-21

Chair of governors

Date:

21-10-21

# Contents

## Statement of Intent

### Preventative measures:

1. Legal framework
2. Ensuring a clean environment
3. Student immunisation
4. Staff immunisation

### In the event of infection:

5. Preventing the spread of infection
6. Vulnerable students
7. Procedures for unwell students/staff
8. Exclusion
9. Medication
10. Outbreaks of infectious diseases
11. Pregnant staff members
12. Staff handling food
13. Managing specific infectious diseases
14. Monitoring and review

### Appendices

- a) **[New]** Infection Control During the Coronavirus Pandemic
- b) **[Updated]** Managing Specific Infectious Diseases
- c) **[Updated]** Infection Absence Periods
- d) Diarrhoea and Vomiting Outbreak Action Checklist
- e) List of Notifiable Diseases

## Statement of Intent

Infections can easily spread in a school due to:

- Students' immature immune systems.
- The close-contact nature of the environment.
- Some students having not yet received full vaccinations.
- Students' poor understanding of good hygiene practices.

Infections commonly spread in the following ways:

- **Respiratory spread** – contact with coughs or other secretions from an infected person.
- **Direct contact spread** – direct contact with the infecting organism, e.g., skin-on-skin contact during sports.
- **Gastrointestinal spread** – contact with contaminated food or water, or contact with infected faeces or unwashed hands.
- **Blood borne virus spread** – contact with infected blood or bodily fluids, e.g., via bites or used needles.

We actively prevent the spread of infection via the following measures:

- Maintaining high standards of personal hygiene and practice
- Maintaining a clean environment
- Routine immunisations
- Taking appropriate action when infection occurs

This policy aims to help school staff prevent and manage infections in school. It is not intended to be used as a tool for diagnosing disease, but rather a series of procedures informing staff what steps to take to prevent infection and what actions to take when infection occurs.

## 1. Legal framework

1.1. This policy has due regard to legislation including, but not limited to, the following:

- Control of Substances Hazardous to Health Regulations 2002 (as amended 2004)
- Health and Safety at Work etc. Act 1974
- The Management of Health and Safety at Work Regulations 1999
- The Reporting of Injuries, Diseases and Dangerous Occurrences Regulations 2013
- The Health Protection (Notification) Regulations 2010

1.2. This policy has due regard to statutory guidance including, but not limited to, the following

- Public Health England (2019) 'Health protection in schools and other childcare facilities'
  - DfE (2015) 'Supporting students at school with medical conditions'
- 1.3. This policy operates in conjunction with the following school policies and documents:
- Health and Safety Policy
  - First Aid Policy

## **2. Ensuring a clean environment**

### **Sanitary facilities**

- 2.1. Wall-mounted soap dispensers are used in all toilets – bar soap is never used.
- 2.2. A foot-operated wastepaper bin is always made available where disposable paper towels are used.
- 2.3. Toilet paper is always available in cubicles.
- 2.4. Suitable sanitary disposal facilities are provided where necessary.

### **Handwashing**

- 2.5. All staff and students are advised to wash their hands after using the toilet, before eating or handling food, and after touching animals.

### **Blood and other bodily fluids**

- 2.6. Cuts and abrasions are covered with waterproof dressings.
- 2.7. When coughing or sneezing, all staff and students are encouraged to cover their nose and mouth with a disposable tissue and dispose of the tissue after use, and to wash their hands afterwards.
- 2.8. Personal protective equipment (PPE) are worn where there is a risk of contamination with blood or bodily fluids during an activity. Gloves are disposable, non-powdered vinyl or latex and CE (Conformité Européene) marked. If there is a risk of splashing to the face, goggles are worn.
- 2.9. Spillages of blood, faeces, saliva, vomit, nasal and eye discharges are cleaned up immediately. They are cleaned using a mixture of detergent and disinfectant. Paper towels or cloths are used, always wearing PPE, and they are disposed of after use.

### **Bites**

- 2.10. If a bite does not break the skin, the affected area is cleaned with soap and water.

- 2.11. If a bite breaks the skin, the affected area is cleaned with soap and running water, the incident is recorded in the student accident log\_and medical advice is sought immediately.

### **3. Student immunisation**

- 3.1. The school keeps up-to-date with national and local immunisation scheduling and advice via [www.nhs.uk/conditions/vaccinations/](http://www.nhs.uk/conditions/vaccinations/).
- 3.2. Each student's immunisation status is checked upon school entry and at the time of any vaccination.
- 3.3. Whilst the school encourages parents to have their children immunised, parental consent will always be sought before a vaccination is given.
- 3.4. The school will ensure that any students with existing medical conditions are medically cleared to be given the vaccine in question.
- 3.5. A healthcare team will visit the school in order to carry out vaccinations and will be able to advise students if there are any concerns.
- 3.6. Girls aged between 12 and 13 can choose to get the HPV vaccine to protect themselves against some types of cervical cancer. This vaccine comprises two injections given 6-12 months apart.
- 3.7. All students aged 14 will be offered the 3-in-1 teenage booster vaccination to top up the effects of the pre-school vaccines against diphtheria, polio and tetanus.
- 3.8. All students aged 14 will be offered the MenACWY vaccine as part of the routine adolescent schools programme.
- 3.9. Any students who become unwell after receiving a vaccination will be treated by the healthcare team who administered the vaccine, following the school's procedures for sick and unwell students.
- 3.10. Any side effects from the vaccinations, such as becoming unwell, will be reported to the healthcare team who administered the vaccination, allowing them to record the symptoms and the time that the vaccine was administered.
- 3.11. Any medication required to relieve the side effects of a vaccination, such as painkillers, will be administered in accordance with the school's Administering Medication Policy.
- 3.12. Regular communication is maintained after students return to lessons, as some side effects can take several hours to develop.
- 3.13. Members of staff will be with students before, during and after vaccinations, in order to keep the students relaxed and create a calming atmosphere.

- 3.14. The school will ensure that the venue used is a clean, open, well-ventilated room, where students can access water and fresh air.
- 3.15. Needles are kept away from students before and after the vaccine is administered.
- 3.16. Some vaccinations may involve an exclusion period in which students are not required to attend school. The administering healthcare team will provide advice in such cases.

#### **4. Staff immunisation**

- 4.1. All staff will undergo a full occupational health check prior to employment, which confirms they are up to date with their immunisations.
- 4.2. Staff should be up to date with immunisations; in particular, we encourage the following:
  - **Hepatitis B:** We do not recommend Hepatitis B vaccines for staff in routine contact with infected children; however, where staff are involved with the care of children with severe learning disabilities or challenging behaviour, we encourage immunisation.
  - **Rubella:** Female staff of childbearing age are encouraged to check with their GP that they are immune to the rubella (German measles) virus. If they are not immune, we encourage them to be immunised with the MMR vaccine, except during pregnancy.

#### **5. Preventing the spread of infection**

Parents should not bring their child to school in the following circumstances:

- The child shows signs of being poorly and needing one-to-one care
- The child has untreated conjunctivitis
- The child has a high temperature/fever
- The child has untreated head lice
- The child has been vomiting and/or had diarrhoea within the last 48 hours
- The child has an infection and the [minimum recommended exclusion period](#) has not yet passed

#### **6. Vulnerable students**

- 6.1. Students with impaired immune defence mechanisms (known as immune-compromised) are more likely to acquire infections. In addition, the effect of an infection is likely to be more significant for such students. These students may have a disease that compromises their immune system or be undergoing treatment, such as chemotherapy, that has a similar effect.

- 6.2. Parents are responsible for notifying the school if their child is “vulnerable”. If a vulnerable child is thought to have been exposed to an infectious disease, the child's parents will be informed and encouraged to seek medical advice from their doctor or specialist.

## **7. Procedures for unwell students/staff**

- 7.1. Staff are required to know the warning signs of students becoming unwell including, but not limited to, the following:

- Not being themselves
- Not having a snack
- Not eating at lunchtimes
- Wanting more attention/sleep than usual
- Displaying physical signs of being unwell, e.g., watery eyes, a flushed face or clammy skin

- 7.2 Where a staff member identifies a student as unwell, the student is taken to main reception where the student's parents will be informed of the situation. Staff will:

- Attempt to cool the student down if they are too hot, by opening a window and suggesting that the student removes their top layers of clothing.
- Provide the student with a drink of water.
- Move the student to a quieter area of the classroom or school.
- Ensure there is a staff member available to comfort the student.
  - Summon emergency medical help if required.

- 7.3 Students and staff displaying any of the of becoming unwell will be sent home, and we will recommend that they see a doctor.

- 7.4 If a student is identified with sickness and diarrhoea, the student's parents will be contacted immediately, and the child will be sent home and may only return after 48 hours have passed without symptoms.

- 7.5 If a staff member is suffering from vomiting and diarrhoea, they will be sent home and may not return until 48 hours have passed without symptoms.

- 7.6 If the school is unable to contact a student's parents in any situation, the student's alternative emergency contacts will be contacted.

### Contaminated clothing

- 7.7 If the clothing of the first-aider or a student becomes contaminated, the clothing is removed as soon as possible and placed in a plastic bag. The student's clothing is sent home with the student, and parents are advised of the best way to launder the clothing.

## 8. Exclusion

- 8.1. Students suffering from infectious diseases will be excluded from school on medical grounds for the [minimum recommended period](#).
- 8.2. Students can be formally excluded on medical grounds by the headteacher.
- 8.3. If parents insist on their child returning to school when the child still poses a risk to others, the LA may serve notice on the child's parents to require them to keep the child away from school until the child no longer poses a risk of infection.
- 8.4. If a student is exposed to an infectious disease, but is not confirmed to be infected, this is not normally a valid reason for exclusion; however, the local health protection team (HPT) may be contacted to advise on a case-by-case basis.

## 9. Medication

- 9.1. Where a student has been prescribed medication by a doctor, dentist, nurse or pharmacist, the first dose will be given at home, in case the student has an adverse reaction.
- 9.2. The student will only be allowed to return to school 24 hours after the first dose of medication, to allow it time to take effect.

## 10. Outbreaks of infectious diseases

- 10.1. An incident is classed as an 'outbreak' where:
  - Two or more people experiencing a similar illness are linked in time or place.
  - A greater than expected rate of infection is present compared with the usual background rate, e.g.:
    - Two or more students in the same classroom are suffering from vomiting and diarrhoea.
    - A greater number of students than usual are diagnosed with scarlet fever.
    - There are two or more cases of measles at the school.
- 10.2. Suspected outbreaks of any of the diseases listed on the [List of Notifiable Diseases](#) will always be reported.
- 10.3. As soon as an outbreak is suspected (even if it cannot be confirmed), the headteacher will contact the HPT to discuss the situation and agree if any actions are needed.
- 10.4. The headteacher will provide the following information:



- The number of staff and children affected
  - The symptoms present
  - The date(s) the symptoms first appeared
  - The number of classes affected
- 10.5. If the headteacher is unsure whether suspected cases of infectious diseases constitute an outbreak, they will contact the HPT.
- 10.6. The HPT will provide the school with draft letters and factsheets to distribute to parents.
- 10.7. The HPT will always treat outbreaks in the strictest confidence; therefore, information provided to parents during an outbreak will never include names and other personal details.
- 10.8. If a student is identified as having a notifiable disease, as outlined in [the guide to Infection Absence Periods](#), the school will inform the parents, who should inform their child's GP. It is a statutory requirement for doctors to then notify their local Public Health England centre.
- 10.9. During an outbreak, enhanced cleaning protocols will be undertaken, following advice provided by the local HPT.

## 11. Pregnant staff members

- 11.1. If a pregnant staff member develops a rash or is in direct contact with someone who has a potentially contagious rash, we will strongly encourage her to speak to her doctor or midwife.
- 11.2. **Chickenpox:** If a pregnant staff member has not already had chickenpox or shingles, becoming infected can affect the pregnancy. If a pregnant staff member believes they have been exposed to chickenpox or shingles and have not had either infection previously, she will speak to her midwife or GP as soon as possible. If a pregnant staff member is unsure whether they are immune, we encourage them to take a blood test.
- 11.3. **Measles:** If a pregnant staff member is exposed to measles, she will inform her midwife immediately. All female staff under the age of 25, who work with young children, are asked to provide evidence of two doses of MMR vaccine or a positive history of measles.
- 11.4. **Rubella (German measles):** If a pregnant staff member is exposed to rubella, she will inform her midwife immediately. All female staff under the age of 25, who work with young children, are asked to provide evidence of two doses of MMR vaccine or a positive history of Rubella.

- 11.5 **Slapped cheek disease (Parvovirus B19):** If a pregnant staff member is exposed to slapped cheek disease, she will inform her midwife promptly.

## **12. Staff handling food**

- 12.1. Food handling staff suffering from transmittable diseases will be excluded from all food handling activity until advised by the local Environmental Health Officer that they are clear to return to work. Both food handling staff and midday assistants are not permitted to attend work if they are suffering from diarrhoea and/or vomiting. They are not permitted to return to work until 48 hours have passed since diarrhoea and/or vomiting occurred, or until advised by the local environmental health officer that they may return to work.
- 12.2 The school will notify the local Environmental Health Department as soon as we are notified that a staff member engaged in the handling of food has become aware that they are suffering from, or likely to be carrying, an infection that may cause food poisoning.
- 12.3. Food handlers are required by law to inform the school if they are suffering from any of the following:
- Typhoid fever
  - Paratyphoid fever
  - Other salmonella infections
  - Dysentery
  - Shigellosis
  - Diarrhoea (where the cause of which has not been established)
  - Infective jaundice
  - Staphylococcal infections likely to cause food poisoning like impetigo, septic skin lesions, exposed infected wounds, boils
  - E. coli VTEC infection
- 12.4. 'Formal' exclusions will be issued where necessary, but employees are expected to provide voluntary 'off work' certificates from their GP.

## **13 Managing specific infectious diseases**

When an infectious disease occurs in the school, we will follow the appropriate procedures set out in the [Managing Specific Infectious Diseases](#) appendix.

## **14 Monitoring and review**

- 14.1. All members of staff are required to familiarise themselves with this policy as part of their induction programme.

- 14.2. The headteacher will review this policy on an annual basis and will make any changes necessary, considering the current effectiveness of infection control and prevention.
- 14.3. The next scheduled review date is June 2021.

## Appendix A

## Infection Control During the Coronavirus Pandemic

### Statement of intent

We understand that we face a time of great uncertainty and, as a school, we are doing all we can to provide clarity and safety for the school community. This appendix includes provisions which the school will have due regard for during the coronavirus pandemic. The information in this section is under constant review and is updated to reflect changes to government guidance as it is released.

### 1. Legal framework

This policy has due regard to all relevant legislation and statutory guidance including, but not limited to, the following:

- DfE and PHE (2020) 'Coronavirus (COVID-19): implementing social distancing in education and childcare settings'
- PHE and DfE (2020) 'Coronavirus (COVID-19): guidance for educational settings'
- PHE (2020) 'COVID-19: cleaning of non-healthcare settings'

### 2. Social distancing measures

2.1. The school will adhere to the government's social distancing guidelines as much as is possible.

2.2. To ensure the risk of virus spread for both staff and students still in school is as low as possible, the following action will be taken:

- Students, parents, carers and any visitors will be told not to attend if they are displaying symptoms of coronavirus.
- The way students arrive at school will be reviewed, with a view to reducing any unnecessary travel on coaches, buses or other public transport.
- Break and lunch time dining room service will be staggered into year group bubbles.
- At lunch time, both students will be asked to queue for meals in year group bubbles and in specified year group areas in the dining room.
- Students will be asked to visit specific year group toilets one after the other.
- Staff will ensure students keep a safe distance.
- All unnecessary staff gatherings will be avoided, e.g. where possible, meetings will take place via video conference from different classrooms or offices.
- Parents will be discouraged from visiting school.

### 3. Additional hygiene and cleaning measures

- 3.1. Anyone who feels unwell will be told to stay at home for 7 days if they live alone, or 14 days if they live with others.
- 3.2 All staff and students will be:
  - Told to frequently wash their hands with soap and water for 20 seconds.
  - Encouraged not to touch their faces.
  - Told to use a tissue or their elbow to catch coughs or sneezes and use bins for tissue waste.
- 3.3. The frequency of cleaning will increase, particularly for surfaces in classrooms, within toilet blocks and in changing rooms, in accordance with PHE's 'COVID19: cleaning of non-healthcare settings' guidance and the school's Cleaning Policy.
- 3.4. Equipment, including keyboards, tables, chairs, door handles, light switches and bannisters, will be cleaned and disinfected regularly.

### 4. Preventing the further spread of infection

- 4.1. If anyone becomes unwell with the symptoms of coronavirus, they will be sent home and advised to follow PHE's 'COVID-19: guidance for households with possible coronavirus infection'.
- 4.2. If a student is awaiting collection, they will be moved to a room where they can be isolated behind a closed door and receive adult supervision, if required. Where possible, a window will be opened for ventilation.
- 4.3. Where isolation is not possible, the student will be moved to an area which is at least two metres from anyone else.
- 4.4. If the student needs to use the toilet whilst waiting to go home, they will be required to use a separate bathroom. This bathroom will be cleaned and disinfected using standard cleaning products before anyone else uses it.
- 4.5. If a member of staff has helped someone who is unwell with coronavirus symptoms, they will not be sent home unless they develop symptoms themselves.
- 4.6. Staff must wash their hands thoroughly for 20 seconds following any contact with someone who is unwell.
- 4.7. When cleaning an area where a person with possible or confirmed coronavirus has been, staff will use disposable gloves and an apron. Staff will wash their hands with soap and water for 20 seconds after all PPE has been removed.
  - 4.8. If there is visible contamination to an area, e.g. with bodily fluids, cleaning staff will use additional PPE to protect their eyes, mouth and nose.

- 4.9. Areas where a symptomatic individual has passed through and spent minimal time, e.g. corridors, but which are not visibly contaminated with bodily fluids, will be cleaned thoroughly with normal cleaning products.
- 4.10. All surfaces that a symptomatic person has come into contact with will be cleaned and disinfected, including objects which are visibly contaminated and those which are potentially contaminated, e.g. door handles.
- 4.11. Disposable cloths or paper rolls and disposable mop heads will be used to clean all hard surfaces, floors, chairs, door handles and sanitary fittings, using one of the following options:
- A combined detergent disinfectant solution at a dilution of 1,000 parts per million available chlorine
  - A household detergent followed by disinfection (1000 ppm av.cl), following the manufacturer's instructions for dilution, application and contact times
  - An alternative disinfectant which is checked to ensure it is effective against enveloped viruses
- 4.12. Cleaning staff will be asked to:
- Avoid creating splashes and spray when cleaning.
  - Dispose of any cloths and mop heads used, by putting them into waste bags.
  - Steam clean items that cannot be cleaned using detergents, e.g. upholstered furniture.
  - Dispose of items that are heavily contaminated with body fluids and cannot be cleaned by washing.
- 4.13. Waste from possible cases and the cleaning of potentially infected areas will be put into a plastic rubbish bag which will be tied when full. The plastic bag will then be placed into a second tied bin bag, put in a suitable and secure place, and marked for storage until the individual's test results are known.

## 5. Communication with parents

The school will inform parents about the measures being taken and request they provide support with implementation, e.g. by encouraging measures to continue at home.

## 6. Students with SEND

- 6.1. The school will work with the LA and parents to decide how best to continue supporting students with EHC plans whilst ensuring they stay healthy and safe.

6.2. Some students with SEND or complex needs may be unable to follow social distancing guidelines. In these circumstances, staff will increase their level of self-protection by minimising close contact (where appropriate), cleaning frequently touched surfaces, and carrying out more frequent handwashing.

## 7. Monitoring and review

7.1. The headteacher is responsible for continually monitoring PHE and DfE updates and updating this appendix in line with any changes to government guidance.

7.2. Any changes to this appendix will be communicated to all staff, parents and relevant stakeholders.

## Appendix B

## Managing Specific Infectious Diseases

Disease	Symptoms	Considerations	Exclusion period
Athlete's foot	Scaling or cracking of the skin, particularly between the toes, or blisters containing fluid. The infection may be itchy.	Cases are advised to see their GP for advice and treatment.	Exclusion is not necessary.
Chicken pox	Sudden onset of fever with a runny nose, cough and generalised rash. The rash then blisters and scabs over. Several blisters may develop at once, so there may be scabs in various stages of development. Some mild infections may not present symptoms.	Cases are advised to consult their GP.	<p>Chickenpox is infectious from 48 hours prior to a rash appearing up to five days after the onset of a rash.</p> <p>Cases will be excluded from school for five days from the onset of a rash.</p> <p>It is not necessary for all the spots to have healed before the case returns to school.</p>
Cold sores	The first signs of cold sores are tingling, burning or itching in the affected area. Around 24 hours after the first signs appear the area will redden and swell, resulting in a fluid-filled blister. After blistering, they break down to form ulcers then dry up and crust over.	Cases are advised not to touch the cold sore, or to break or pick the blisters. Sufferers of cold sores should avoid kissing people and should not share items such as cups, towels and facecloths.	Exclusion is not necessary.
Conjunctivitis	The eye(s) become reddened and swollen, and there may be a yellow or green discharge. Eyes may feel itchy and 'gritty'.	<p>Cases are encouraged to seek advice, wash their hands frequently and not to rub their eyes.</p> <p>The HPT will be contacted if an outbreak occurs.</p>	Exclusion is not necessary.



Disease	Symptoms	Considerations	Exclusion period
Coronavirus	A new, continuous cough and a high temperature are the main symptoms of coronavirus.	Cases will be sent home and advised to contact 111 for advice, or 999 if they become seriously ill or believe their life is at risk.  The local HPT will be contacted if an outbreak occurs.	Cases will be required to self-isolate at home for 7 days if they live alone, or 14 days if they live with others.
Food poisoning	Symptoms normally appear within one to two days of contaminated food being consumed, although they may start at any point between a few hours and several weeks later. The main symptoms are likely to be nausea, vomiting, diarrhoea, stomach cramps and fever.	Cases will be sent home.  The HPT will be contacted where two or more cases with similar symptoms are reported.  The cause of a food poisoning outbreak will always be investigated.	Cases will be excluded until 48 hours have passed since symptoms were present.  For some infections, longer exclusion periods may be required. The HPT will advise in such cases.
Giardia	Symptoms include abdominal pain, bloating, fatigue and pale, loose stools.	Cases will be sent home.  The HPT will be contacted where two or more cases with similar symptoms are reported.	Cases will be excluded until 48 hours have passed since symptoms were present.
Salmonella	Symptoms include diarrhoea, headache, fever and, in some cases, vomiting.	Cases will be sent home.  The HPT will be contacted where two or more cases with similar symptoms are reported.	Cases will be excluded until 48 hours have passed since symptoms were present.

Disease	Symptoms	Considerations	Exclusion period
Typhoid and paratyphoid fever	Symptoms include tiredness, fever and constipation. The symptoms of paratyphoid fever include fever, diarrhoea and vomiting.	All cases will be immediately reported to the HPT.	<p>Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved.</p> <p>Environmental health officers or the HPT may advise the school to issue a lengthened exclusion period.</p>
E. coli (verocytotoxigenic or VTEC)	Symptoms vary but include diarrhoea, abdominal cramps, headaches and bloody diarrhoea.	Cases will immediately be sent home and advised to speak to their GP.	<p>Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved.</p> <p>Where the sufferer poses an increased risk, for example, food handlers, they will be excluded until a negative stool sample has been confirmed.</p> <p>The HPT will be consulted in all cases.</p>
Gastroenteritis	Symptoms include three or more liquid or semiliquid stools in a 24-hour period.	The HPT will be contacted where there are more cases than usual.	<p>Cases will be excluded until 48 hours have passed since symptoms were present. If medication is prescribed, the full course must be completed and there must be no further symptoms displayed for 48 hours following completion of the course before the cases may return to school.</p> <p>Cases will be excluded from swimming for two weeks following their last episode of diarrhoea.</p>

Disease	Symptoms	Considerations	Exclusion period
Bacillary dysentery (Shigella)	Symptoms include bloody diarrhoea, vomiting, abdominal pain and fever. It lasts four to seven days on average, but potentially several weeks.	The school will contact the HPT.	Microbiological clearance is required for some types of shigella. The HPT will advise.
Campylobacter	Symptoms include diarrhoea, headache, fever and, in some cases, vomiting.		Cases will be excluded until 48 hours have passed since symptoms were present.
Cryptosporidiosis	Symptoms include abdominal pain, diarrhoea and occasional vomiting.		Cases will be excluded until 48 hours have passed since symptoms were present.
Glandular fever	Symptoms include severe tiredness, aching muscles, sore throat, fever, swollen glands and occasionally jaundice.	The sufferer may feel unwell for several months and the school will provide reasonable adjustments where necessary.	Exclusion is not necessary, and cases can return to school as soon as they feel well.
Hand, foot and mouth disease	Symptoms include a fever, reduced appetite and generally feeling unwell. One or two days later, a rash with blisters will develop on cheeks, hands and feet. Not all cases will have symptoms.		Exclusion is not necessary, and cases can return to school as soon as they feel well.
Head lice	Other than the detection of live lice or nits, there are no immediate symptoms until two to three weeks after infection, where itching and scratching of the scalp occurs.	Treatment is only necessary when live lice are seen.  Staff are not permitted to inspect any student's hair for head lice.	Exclusion is not necessary.

Disease	Symptoms	Considerations	Exclusion period
		<p>If a staff member incidentally notices head lice in a student's hair, they will inform the student's parents and advise them to treat their child's hair.</p> <p>When a student has been identified as having a case of head lice, a letter will be sent home to all parents notifying them that a case of head lice has been reported and asking all parents to check their children's hair.</p>	
Hepatitis A	Symptoms include abdominal pain, loss of appetite, nausea, fever and tiredness, followed by jaundice, dark urine and pale faeces.	The illness in children usually lasts one to two weeks but can last longer and be more severe in adults.	<p>Cases are excluded while unwell and for seven days after the onset of jaundice (or the onset of symptoms if no jaundice presents), the case is under five years of age or where hygiene is poor.</p> <p>There is no need to exclude older children with good hygiene.</p>
Hepatitis B	Symptoms include general tiredness, nausea, vomiting, loss of appetite, fever and dark urine, and older cases may develop jaundice.	<p>The HPT will be contacted where advice is required.</p> <p>The procedures for dealing with blood and other bodily fluids will always be followed.</p> <p>The accident book will always be completed with details of injuries or adverse events related to cases.</p>	<p>Acute cases will be too ill to attend school and their doctor will advise when they are fit to return.</p> <p>Chronic cases will not be excluded or have their activities restricted.</p> <p>Staff with chronic hepatitis B infections will not be excluded.</p>

Disease	Symptoms	Considerations	Exclusion period
Hepatitis C	Symptoms are often vague but may include loss of appetite, fatigue, nausea and abdominal pain. Less commonly, jaundice may occur.	The procedures for dealing with blood and other bodily fluids will always be followed.  The accident book will always be completed with details of injuries or adverse events related to cases.	Cases will not be excluded or have their activities restricted.
Impetigo	Symptoms include lesions on the face, flexures and limbs.	Towels, facecloths and eating utensils will not be shared by students.  Toys and play equipment will be cleaned thoroughly.	Cases will be excluded until lesions have healed and crusted or 48 hours after commencing antibiotic treatment.
Influenza	Symptoms include headache, fever, cough, sore throat, aching muscles and joints, and tiredness.	Those in risk groups will be encouraged to have the influenza vaccine.  Anyone with flu-like symptoms will stay home until they have recovered. Students under 16 will not be given aspirin.	Cases will remain home until they have fully recovered.
Measles	Symptoms include a runny nose, cough, conjunctivitis, high fever and small white spots around the cheeks. Around the third day, a rash of flat red or brown blotches may appear on the face then spread around the body.	All students are encouraged to have MMR immunisations in line with the national schedule.  Staff members should be up to date with their MMR vaccinations.  Pregnant staff members and those with weak immune systems will be encouraged to contact their GP immediately for advice if they come into contact with measles.	Cases are excluded for four days after the onset of a rash.

Disease	Symptoms	Considerations	Exclusion period
Meningitis	Symptoms include fever, severe headaches, photophobia, stiff neck, non-blanching rash, vomiting and drowsiness.	Meningitis is a notifiable disease.	Once a case has received any necessary treatment, they can return to school.
Meningococcal meningitis and meningitis septicaemia	Symptoms include fever, severe headaches, photophobia, stiff neck and a non-blanching rash.	<p>Medical advice will be sought immediately.</p> <p>The confidentiality of the case will always be respected.</p> <p>The HPT and school health advisor will be notified of a case of meningococcal disease in the school. The HPT will conduct a risk assessment and organise antibiotics for household and close contacts.</p> <p>The HPT will be notified if two cases of meningococcal disease occur in the school within four weeks.</p>	<p>When the case has been treated and recovered, they can return to school.</p> <p>Exclusion is not necessary for household or close contacts unless they have symptoms suggestive of meningococcal infection.</p>
Meningitis (viral)	Symptoms include headache, fever, gastrointestinal or upper respiratory tract involvement and, in some cases, a rash.	<p>The case will be encouraged to consult their GP.</p> <p>If more than once case occurs, the HPT will be consulted.</p>	No exclusion is required.

Disease	Symptoms	Considerations	Exclusion period
Meticillin resistant staphylococcus aureus (MRSA)	Symptoms are rare but include skin infections and boils.	All infected wounds will be covered.	No exclusion is required.
Mumps	Symptoms include a raised temperature and general malaise. Then, stiffness or pain in the jaws and neck is common. Following this, the glands in the cheeks and under the jaw swell up and cause pain (this can be on one or both sides). Mumps may also cause swelling of the testicles.	The case will be encouraged to consult their GP.  Parents are encouraged to immunise their children against mumps.	Cases can return to school five days after the onset of swelling if they feel able to do so.
Ringworm	Symptoms vary depending on the area of the body affected.	Students with ringworm of the feet will always wear socks and trainers and cover their feet during physical education.	No exclusion is usually necessary.  For infections of the skin and scalp, cases can return to school once they have received treatment.
Rotavirus	Symptoms include severe diarrhoea, stomach cramps, vomiting, dehydration and mild fever.	Cases will be sent home if unwell and encouraged to speak to their GP.	Cases will be excluded until 48 hours have passed since symptoms were present.
Rubella (German Measles)	Symptoms are usually mild, with a rash being the first indication. There may also be mild catarrh, headaches or vomiting. There may be a slight fever and some tenderness in the neck, armpits or groin, and there may be joint pains.	MMR vaccines are promoted to all students.	Cases will be excluded for six days from the appearance of the rash.
Scabies	Symptoms include tiny pimples and nodules on a rash, with burrows commonly seen on the wrists, palms, elbows, genitalia and buttocks.	All household contacts and any other very close contacts should have one treatment at the same time as the second treatment of the case.	Cases will be excluded until after the first treatment has been carried out.

Disease	Symptoms	Considerations	Exclusion period
		The second treatment must not be missed and should be carried out one week after the first treatment.	
Scarlet Fever	Symptoms include acute inflammation of the pharynx or tonsils, with tonsils reddening in colour and becoming partially covered with a thick, yellowish exudate. In severe cases, there may be a high fever, difficulty swallowing and tender, enlarged lymph nodes. A rash develops on the first day of fever and is red, generalised, pinhead in size and gives the skin a sandpaper-like texture, with the tongue developing a strawberry like appearance.	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered.  If two or more cases occur, the HPT will be contacted.	Cases are excluded for 24 hours following appropriate antibiotic treatment.
Slapped cheek syndrome, Parvovirus B19, Fifth's Disease	Where symptoms develop, they include a rosered rash making the cheeks appear bright red.	Cases will be encouraged to visit their GP.	Exclusion is not required.
Threadworm	Symptoms include itching around the anus, particularly at night.	Cases will be encouraged to visit their GP.	Exclusion is not required.
Tuberculosis (TB)	Symptoms include cough, loss of appetite, weight loss, fever, sweating (particularly at night), breathlessness and pains in the chest. TB in parts of the body other than the lungs may produce a painful lump or swelling.	Advice will be sought from the HPT before taking any action, and regarding exclusion periods.	Cases with infectious TB can return to school after two weeks of treatment if well enough to do so and provided they have responded to anti-TB therapy. Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses,



Disease	Symptoms	Considerations	Exclusion period
			will not be excluded.
Whooping cough (pertussis)	Symptoms include a heavy cold with a persistent cough. The cough generally worsens and develops the characteristic 'whoop'. Coughing spasms may be worse at night and may be associated with vomiting.	Cases will be advised to see their GP. Parents are advised to have their children immunised against whooping cough.	Cases will not return to school until they have had 48 hours of appropriate treatment with antibiotics and feel well enough to do so, or 21 days from the onset of illness if no antibiotic treatment is given. Cases will be allowed to return in the above circumstances, even if they are still coughing.

## Appendix C                      Infection Absence Periods

This table details the minimum required period for staff and students to stay away from school following an infection, as recommended by Public Health England.

- \* Identifies a notifiable disease. It is a statutory requirement that doctors report these diseases to their local Public Health England centre.

Infection	Recommended minimum period to stay away from school	Comments
Athlete's foot	None	Treatment is recommended; however, this is not a serious condition.
Chicken pox	Until all vesicles have crusted over	Follow procedures for vulnerable children and pregnant staff.
Cold sores	None	Avoid contact with the sores.
Conjunctivitis	None	If an outbreak occurs, consult the HPT.
Coronavirus	Until fully recovered and no other member of the same household is presenting symptoms (7 days if living alone, 14 days if living with others)	If coronavirus is suspected, consult the local HPT.
Diarrhoea and/or vomiting	Whilst symptomatic and 48 hours from the last episode	GPs should be contacted if diarrhoea or vomiting occur after taking part in water-based activities.
Diphtheria*	Exclusion is essential.	Family contacts must be excluded until cleared by the HPT and the HPT must always be consulted.

Infection	Recommended minimum period to stay away from school	Comments
Flu (influenza)	Until recovered	Report outbreaks to the HPT.
Glandular fever	None	
Hand foot and mouth	None	Contact the HPT if a large number of children are affected. Exclusion may be considered in some circumstances.
Head lice	None	Treatment recommended only when live lice seen.
Hepatitis A*	Seven days after onset of jaundice or other symptoms	If it is an outbreak, the HPT will advise on control measures.
Hepatitis B*, C* and HIV	None	Not infectious through casual contact. Procedures for bodily fluid spills must be followed.
Impetigo	48 hours after commencing antibiotic treatment, or when lesions are crusted and healed	Antibiotic treatment is recommended to speed healing and reduce the infectious period.
Measles*	Four days from onset of rash	Preventable by vaccination (MMR). Follow procedures for vulnerable children and pregnant staff.
Meningococcal meningitis*/septicaemia*	Until recovered	Meningitis ACWY and B are preventable by vaccination. The HPT will advise on any action needed.

Infection	Recommended minimum period to stay away from school	Comments
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. The HPT will advise on any action needed.
Meningitis viral*	None	As this is a milder form of meningitis, there is no reason to exclude those who have been in close contact with infected persons.
MRSA	None	Good hygiene, in particular environmental cleaning and handwashing is important to minimise the spread. The local HPT should be consulted.
Mumps*	Five days after onset of swelling	Preventable by vaccination with two doses of MMR.
Ringworm	Exclusion is not usually required	Treatment is required.
Rubella (German measles)	Four days from onset of rash	Preventable by two doses of immunisation (MMR). Follow procedures for pregnant staff.
Scarlet fever	24 hours after commencing antibiotic treatment	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered. If two or more cases occur, the HPT should be contacted.
Scabies	Can return to school after first treatment	The infected person's household and those who have been in close contact will also require treatment.
Slapped cheek/Fifth disease/Parvo Virus B19	None (once rash has developed)	Follow procedures for vulnerable children and pregnant staff.
Threadworms	None	Treatment recommended for the infected person and household contacts.
Tonsillitis	None	There are many causes, but most causes are virus-based and do not require antibiotics.

Infection	Recommended minimum period to stay away from school	Comments
Whooping cough (pertussis)*	Two days from commencing antibiotic treatment, or 21 days from the onset of illness if no antibiotic treatment is given	Preventable by vaccination. Non-infectious coughing can continue for many weeks after treatment. The HPT will organise any necessary contact tracing.
Tuberculosis (TB)	Students with infectious TB can return to school after two weeks of treatment if well enough to do so, and as long as they have responded to anti-TB therapy.	Only pulmonary (lung) TB is infectious. It requires prolonged close contact to spread. Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses, should not be excluded. Consult the local HPT before disseminating information to staff and parents.
Warts and verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms.

## Appendix D

## Diarrhoea and Vomiting Outbreak Action Checklist

<b>Date:</b>	
<b>Completed by:</b>	

Action	Action taken?		Comments
	Yes	No	
A 48-hour exclusion rule has been enforced.			
Liquid soap and paper hand towels are available.			
Enhanced cleaning is undertaken twice daily, and an appropriate disinfectant is used.			
Appropriate personal protective equipment (PPE) is available.			
Appropriate waste disposal systems are available for removing infectious waste.			
Toys are cleaned and disinfected on a daily basis.			
Infected linen is segregated, and dissolvable laundry bags are used where possible.			
Visitors are restricted, and essential visitors are informed of the outbreak.			
New children joining the school are delayed from joining.			
The health protection team (HPT) has been informed of any infected food handlers.			
Staff work in dedicated areas and food handling is restricted.			
All staff (including agency) are asked if they are unwell.			
Staff are restricted from working elsewhere.			
The HPT is informed of any planned events at the school.			

## **Appendix E            List of Notifiable Diseases**

Under the Health Protection (Notification) Regulations 2010, the following diseases will always be reported to the health protection team (HPT):

- Acute encephalitis
- Acute meningitis
- Acute poliomyelitis
- Acute infectious hepatitis
- Anthrax
- Botulism
- Brucellosis
- Cholera
- COVID-19
- Diphtheria
- Enteric fever (typhoid or paratyphoid fever)
- Food poisoning
- Haemolytic uremic syndrome (HUS)
- Infectious bloody diarrhoea
- Invasive group A streptococcal disease and scarlet fever
- Legionnaires' disease
- Leprosy
- Malaria
- Measles
- Meningococcal septicaemia
- Mumps
- Plague
- Rabies
- Rubella
- SARS
- Smallpox
- Tetanus
- Tuberculosis
- Typhus
- Viral haemorrhagic fever (VHF)
- Whooping cough
- Yellow fever